

FOR MORE INFORMATION

For more information on this subject, write to the Office of Cancer Communications, National Center Institute, Bethesda, MD 20892
Or call the toll-free number of the Cancer Information Service at **1-800-4-CANCER** OR
CLICK ON THE Virginia Urology Center at
<http://www.uro.com/tcancer.htm>

The University of Wisconsin-Whitewater is committed to equal opportunity in its education programs, activities and employment policies for all persons regardless of race, color, gender, creed, religion, age, ancestry, national origin, disability, sexual orientation, political affiliation, marital status, Vietnamese veteran status, parental status, and pregnancy.

Document Name: T:\UHCS\CATALOG\Brochures\Wellness-Prevention\TSEexam.doc
Review Date: 02/15/05
Next Review: 02/15/06

UNIVERSITY HEALTH AND COUNSELING SERVICES

www.uww.edu/stdrscs/uhcs
Corner of Prairie and Starin
Division of Student Affairs, UW-Whitewater
www.uww.edu/staffair

Hours

Monday-Friday 8:00 - 4:30; summer & evening hours may vary.

Health Services 472-1300

- * Examinations and treatment for illnesses and minor injuries
- * Laboratory tests and medications
- * Allergy injections and immunizations
- * Pelvic exam and Pap test
- * Contraceptives
- * STD and HIV antibody testing
- * Cold self care
- * Referrals to specialists
- * Phone consultations with RN
- * Rehabilitation for students with disabilities

Counseling Services 472-1305

- * Individual counseling for many concerns, including...
 - * Relationships
 - * Depression
 - * Self-esteem
 - * Values clarification
 - * Alcohol and other drugs use
 - * Sexual assault and abuse
 - * Decision-making
 - * Anxiety
 - * Sexual orientation
 - * Stress
- * Group counseling options
- * Consultations with staff
- * Referrals to community agencies
- * Crisis intervention
- * Practicum training site

Wellness Services 472-1300 ext. 2244

- * Lobby Resource Center
- * Royal Purple column
- * Bulletin board kits
- * Educational presentations
- * Web site
- * Posters
- * Wellness booths
- * Program kits

Employee Assistance Program 472-1305

- * Confidential assistance and referral for employees with concerns such as:
 - * Stress
 - * Depression
 - * Finances
 - * Alcohol and other drug use
 - * Troubled coworkers
 - * Marriage/family
 - * Health

24-Hour Emergency help:

911 Whitewater Rescue Squad

472-1060 Sexual Assault Response Team
262-741-3200 or 1-800-365-1587 Crisis Line

Testicular Self Exam

Specializing in the Unique Needs of College Students

Testicular cancer is the most common type of cancer in men ages 20 to 35. Yet, because it accounts for only about percent of all cancers in men, many people have never heard of this type of cancer.


Testicular cancer is of special concern to young men. It can occur anytime after age 15. It is less common in middle-aged and older men. White men are four times more likely to develop testicular cancer than black men. The rate among Hispanic men lies between those of blacks and whites.

Two groups of men have a greater risk of developing testicular cancer – those whose testicles have not descended into the scrotum and those whose testicles descended after age 6. Testicular cancer is 3 to 17 times more likely to develop in these men.

Testicles are male reproductive organs. They produce and store sperm. They also produce testosterone, a hormone that causes such male traits as facial hair and lower voice pitch. Testicles are smooth, oval-shaped, and somewhat firm to the touch. They are below the penis in a sac of skin called the scrotum.

The testicles normally descend into the scrotum before birth. Parents should have their infant sons examined by a doctor to be sure that the testicles have properly descended. If they have not, this can be easily corrected with surgery.

Fifteen years ago, testicular cancer was often fatal because it spread quickly to vital organs such as the lungs. Today, due to advances in treatment, testicular cancer is one of the most curable cancers, especially if detected and treated promptly.



SYMPTOMS

The most common symptom of testicular cancer is a small, painless lump in a testicle or a slightly enlarged testicle. It is important for men to become familiar with the size and feeling of their normal testicles, so that they can detect changes if they occur.

Other possible symptoms include a feeling of heaviness in the scrotum, a dull ache in the lower stomach or groin, a change in the way a testicle feels, or a sudden accumulation of blood or fluid in the scrotum. These symptoms can also be caused by infections or other conditions that are not cancer. A doctor can tell you if you have cancer and what the proper treatment should be.



HOW TO DO TSE

A simple procedure called testicular self-exam (TSE) can increase the chances of finding a tumor early.

Men should perform TSE once a month – after a warm bath or shower. The heat causes the scrotal muscle to relax, making it easier to find anything unusual. TSE is simple and only takes a few minutes.

- ◆ Examine each testicle gently with both hands. The index and middle fingers should be placed underneath the testicle while the thumbs are placed on top. Roll the testicle gently between the thumbs and fingers. One testicle may be larger than the other. This is normal.
- ◆ The epididymis is a cord-like structure on the top and back of the testicle that stores and transports the sperm. Do not confuse the epididymis with an abnormal lump.
- ◆ Feel for any abnormal lumps – about the size of a pea – on the front or side of the testicle. These lumps are usually painless.

If you find a lump, you should contact your doctor right away. The lump may be due to an infection, and a doctor can decide the proper treatment. If the lump is not an infection, it is likely to be cancer. Remember that testicular cancer is highly curable, especially when detected and treated early.

Testicular cancer almost always occurs in only one testicle, and the other testicle is all that is needed for full sexual function.

Routine testicular self-exams are important, but they cannot substitute for doctor's examination. Your doctor should examine your testicles when you have a physical exam. You also can ask your doctor to check the way you do TSE.